

N O T I C E

THIS DOCUMENT HAS BEEN REPRODUCED FROM
MICROFICHE. ALTHOUGH IT IS RECOGNIZED THAT
CERTAIN PORTIONS ARE ILLEGIBLE, IT IS BEING RELEASED
IN THE INTEREST OF MAKING AVAILABLE AS MUCH
INFORMATION AS POSSIBLE

"Made available under NASA sponsorship
in the interest of early and wide dis-
semination of Earth Resources Survey
Program information and without liability
for any use made thereof."

80-10155
CR-143172

SDSU-RSI-80-03

HCMM ENERGY BUDGET DATA AS A MODEL INPUT FOR ASSESSING
REGIONS OF HIGH POTENTIAL GROUNDWATER POLLUTION

(E80-10155) HCMM ENERGY BUDGET DATA AS A	N80-26738
MODEL INPUT FOR ASSESSING REGIONS OF HIGH	
POTENTIAL GROUNDWATER POLLUTION Progress	
Report, Jan. - Mar. 1980 (South Dakota State	Unclass
Univ.) 7 p EC A02/MF A01	CSCL 08H G3/43 00155

Principal Investigator: Donald G. Moore

Report Author: J.L. Heilman

March 1980

Interim Type II Report for January-March, 1980

Prepared for:

Goddard Space Flight Center
Greenbelt, Maryland 20771

RECEIVED

MAY 27 1980

SIS/902.6
HCM 032
TYPE II

TECHNICAL REPORT STANDARD TITLE PAGE

1. Report No. SDSU-RSI-80-03		2. Government Accession No.		3. Recipient's Catalog No.	
4. Title and Subtitle HCMM Energy Budget Data as a Model Input for Assessing Regions of High Potential Groundwater Pollution.				5. Report Date	
				6. Performing Organization Code	
7. Author(s) J.L. Heilman				8. Performing Organization Report No.	
9. Performing Organization Name and Address Remote Sensing Institute South Dakota State University Brookings, South Dakota 57007				10. Work Unit No.	
				11. Contract or Grant No. NAS5-24206	
12. Sponsoring Agency Name and Address NASA Goddard Space Flight Center Greenbelt, MD 20771				13. Type of Report and Period Covered Type II January-March, 1980	
				14. Sponsoring Agency Code	
15. Supplementary Notes					
16. Abstract Significant relationships have been found between surface soil temperatures estimated from HCMM radiometric temperatures and depth to groundwater and near surface soil moisture.					
17. Key Words Suggested by Author Groundwater, thermal inertia, heat sink, HCMM				18. Distribution Statement	
19. Security Classif. (of this report) Unclassified		20. Security Classif. (of this page) Unclassified		21. No. of Pages	
				22. Price	

A. Problems

None

B. Accomplishments

Analyses of day-night registered and apparent thermal inertia data have begun.

C. Significant Results

Previous progress reports (SDSU-RSI-79-01; SDSU-RSI-80-01) documented results relating to use of aircraft thermal data for evaluating depth to groundwater and soil moisture. These results are now being extended to HCMM data.

1. Groundwater

September 4, 1978, HCMM day thermal data were analyzed to assess depth to groundwater in the test site. HCMM apparent temperature were corrected for atmospheric effects using lake temperature of the Oahe Reservoir in central South Dakota. Surface soil temperatures were estimated using the equation

$$T_s = 0.79 T_{HCMM} \times e^{(-0.80 PC)} + 20.35 \quad (1)$$

where T_s is surface soil temperature, T_{HCMM} 's radiometric temperature from HCMM, and PC is percent cover expressed as a fraction. This equation was developed from ground studies and was presented in earlier reports.

A significant relationship was found between T_s and depth to groundwater (Fig. 1). Depth to groundwater was correlated with soil water content (Fig. 2). Thus, the change in T_s shown in Fig. 1 was probably due to both the heat sink effects of the groundwater and thermal inertia effects




Fig. 1. Relationship of predicted soil surface temperatures to depth to groundwater.




Fig. 2. Soil water content in the 0 to 50 cm layer as a function of depth to groundwater.

of soil moisture.

2. Soil Moisture

Three dates of HCMM data have been analyzed for soil moisture. A significant relationship was found between the surface soil-maximum air temperature differential and soil water content (% of field capacity) in the 0 and 4 cm layer of the profile (Fig. 3). T_s was predicted using HCMM radiometric temperature and equation (1). Land use for the data points in Fig. 3 consisted of row crops, small grains, stubble, and pasture. These results demonstrate a potential for empirically relating HCMM data to soil water content for a variety of land uses.

Groundwater and soil moisture analyses are continuing with other dates of HCMM data.

D. Publications

None

E. Recommendations

None

F. Funds Expended

\$83,453.85




Fig. 3. Relationship of predicted soil surface temperature to soil water content in the 0 to 4 cm layer of the profile.